



## Research Article

## Section: Medicine

### Scoring Systems Using Bedside Index of Severity in Acute Pancreatitis (BISAP) and PANC3 Vs Revised Atlanta Classification in Predicting the Severity of Acute Pancreatitis

Dr. Anubha Garg<sup>1</sup>, Dr. Mahima Khatkar<sup>2</sup>, Dr. Bharti<sup>3</sup>, Dr. Dinesh Kumar Garg<sup>4</sup> & Dr. Sumit Chawla<sup>\*5</sup>

<sup>1</sup> Professor, Department of Medicine, Pt. BDS PGIMS, Rohtak.

<sup>2</sup> Junior Resident, Department of Medicine, PGIMS, Rohtak

<sup>3</sup> Associate Professor, Department of Community Medicine, SABVGM, Chhainsa, Faridabad

<sup>4</sup> ASMO, Rohtak, HCMS, Haryana.

<sup>5</sup> Associate Professor, Department of Community Medicine, SABVGM, Chhainsa, Faridabad

## ARTICLE INFO

### Article History:

Received: 05-04-2025

Accepted: 14-05-2025

### Keywords:

PANC3

Pancreatitis

Score

### \*Corresponding author:

Dr. Sumit Chawla, Associate  
Professor, Department of Community  
Medicine, SABVGM, Chhainsa,  
Faridabad

## ABSTRACT

**Background:** Acute pancreatitis is a disease of great importance in clinical practice, defined as acute inflammatory process of the pancreas that may involve local tissues and affect other organs and requires intensive care. **Methods:** This prospective comparative observational study included patients with acute pancreatitis. Bedside index of severity in acute pancreatitis (BISAP) and PANC3 were used to stratify through severity of disease. Score from each model was compared to clinical severity defined by Revised Atlanta classification 2021 for predicting complication and mortality. Sensitivity, specificity and accuracy were compared for each model. A p value of < 0.05 was significant. **Results:** It was seen that, 48 (82.8%) patients had mild to moderately severe acute pancreatitis, while 10 (17.2%) patients had severe acute pancreatitis. BISAP score had a higher sensitivity (88.24%) compared to PANC3 score (76.47%) whereas both had equal specificity for prediction of complications (95.12%). BISAP scoring model had a higher accuracy of 93% for predicting local complications as compared to PANC3 score (86%). **Conclusions:** BISAP score had more sensitivity, diagnostic accuracy, positive and negative predictive value, and equal specificity in predicting the severity of acute pancreatitis as compared to the PANC3 score. Hence, BISAP score was found to predict more number of patients with, the likelihood of progressing to severe disease.

## INTRODUCTION

Acute pancreatitis is as an acute inflammatory process of the pancreas due to activation of digestive enzymes. Acute pancreatitis is defined as by a patients meeting two of the following three criteria [1] Symptoms ( e.g. acute onset epigastric and/or left upper quadrant pain, often radiating to the back), [2] A serum amylase or lipase level greater than 3 times the upper limit of the laboratory reference range, and [3] Radiologic imaging consistent with pancreatitis, usually using CT or MRI.

The incidence and global distribution of acute pancreatitis is heterogenous due to its self limiting nature in mild cases as well as variation in health care systems across different countries. The prevalence rate for pancreatitis in India is 7.9 per 1,00,000. The prevalence rate for men and women is 8.6 and 8.0 per 1,00,000 in India<sup>1</sup>. In India gall stone and alcohol are the most common cause of acute pancreatitis[2].

The severity of organ failure caused by acute pancreatitis is the most important determinant of mortality in the disease. Over 80% of patients have mild self limiting disease while approximately 20% of patients have a severe form with mortality rate as high as 30%[3]. Therefore it is of foremost importance to assess the severity and identify the patients who are at risk for the development of persistent organ failure early in the course of disease so that timely intensive therapy and appropriate interventions are ensured to decrease the mortality rate.

Several scoring systems such as Revised Atlanta classification, Ranson's Criteria, The Acute Physiology and Chronic Health Evaluation II score (APACHE II), The Glasgow score and Harmless Acute Pancreatitis (HAP) Score have been validated and used for assessing the severity of acute pancreatitis. The Atlanta Classification has been considered the global standard tool for the assessment of acute pancreatitis severity.[4] However due to their

complexity, simple scoring systems such as Bedside index of severity in acute pancreatitis (BISAP) and PANC3 score were developed and evaluated. BISAP and PANC3 are simple and can be calculated bedside at the time of admission, requiring less time and fewer laboratory parameters. In this background we are comparing the scoring systems of BISAP and PANC3 vs Revised atlanta classification in predicting the severity of acute pancreatitis. Although various scoring models exist to clinically evaluate the severity of acute pancreatitis and organ failure hitherto, no single system has been considered ideal and there is no consensus on which scoring system to be used. In this context, the need for an objective way to predict acute pancreatitis severity remains an enigma.

## METHODS

This prospective comparative observational study was conducted from March 2021 to March 2022. A total of 58 cases as per diagnostic criteria of acute pancreatitis admitted in the medicine ward of Pt. B.D. Sharma PGIMS Rohtak were enrolled in the study after informed consent. Patients aged 18 years and above, diagnosed of AP (either first attack or recurrent attacks), presenting with acute onset of persistent severe epigastric pain, with or without radiation, and increased serum amylase and lipase levels were included. However, Patients with pre-existing chronic pancreatitis, Chronic Cardiac, liver, lung and kidney disease were excluded from the study.

### Assessment of severity and associated complications:

BISAP, PANC3 and Revised atlanta classification

(RAC) were used to stratify the severity of disease. All subjects thereafter were followed up for 4 weeks to watch for any complications associated with pancreatitis and mortality.

The components of BISAP scoring system are:

- I. Blood Urea Nitrogen (BUN) >25
- II. Impaired mental status
- III. Severe Inflammatory Response Syndrome (SIRS)  $\geq 2$
- IV. Age >60
- V. Pleural effusion

A score >2 indicates severe pancreatitis. SIRS includes two or more of the following conditions: Temperature >38.3°C or <36.0°C, Heart rate of >90 beats/minute, Respiratory rate of >20 breaths/minute or PaCO<sub>2</sub> of <32 mmHg, WBC count of >12,000 cells/mm<sup>3</sup>, <4000 cells/mm<sup>3</sup>, or >10 % immature (band) forms.

The components of PANC3 score are:

- i. Serum hematocrit >44%
- ii. Body Mass Index (BMI) >32 kg/m<sup>2</sup>
- iii. Pleural effusion on the chest x ray

Revised Atlanta classification (RAC) of acute pancreatitis<sup>5</sup> was classified into Mild acute pancreatitis (No organ failure and no local or systemic complications), Moderately severe acute pancreatitis (Organ failure that resolves within 48 h (transient organ failure) and/or Local or systemic complications without persistent organ failure and Severe acute pancreatitis [Persistent organ failure (>48 hours)]. The modified Marshall scoring system is used in the Revised Atlanta classification, and it scores the respiratory (PaO<sub>2</sub>/FiO<sub>2</sub>), cardiovascular (SBP) and renal system (S. creatinine value) to detect organ failure with a score  $\geq 2$  indicating organ failure.


**Table 1: Type of pancreatitis in study subjects (n=58)**

| Type of pancreatitis | No. | %    |
|----------------------|-----|------|
| Mild                 | 41  | 70.7 |
| Moderate             | 7   | 12.1 |
| Severe               | 10  | 17.2 |


In our study, it was also seen that, 48 (82.8%) patients had mild to moderately severe acute pancreatitis, while 10 (17.2%) patients had severe acute pancreatitis which was comparable to other study [8] as shown in table 1.

Most common complication in our study was Acute Kidney Injury (AKI) as shown in figure 1. BISAP score had a higher sensitivity (88.24%) compared to PANC3 score (76.47%) whereas both had equal specificity for prediction

prediction of complications (95.12%). BISAP scoring model had a higher accuracy of 93% for predicting local complications as compared to PANC3 score (86%) (Figure 2). Results of BISAP score were comparable with study done by Lalithkumar et al. [9] which showed that BISAP score had better specificity (95.35%), and diagnostic accuracy (92%) and Park et al [10] which showed that BISAP sensitivity for organ failure was 91.3% and, specificity was 85%.




**Figure 1: Complications in pancreatitis subjects (n=58)**




**Figure 2: Diagnostic value of BISAP and PANC3 score for prediction of complications in pancreatitis subjects**

Systemic inflammatory response syndrome (SIRS) score study done by Alaarabiou A[11] ( 50%) and Vikesh K of 2 or more on day 1 was present in 50 percent of all acute pancreatitis subjects which were similar to results of other



**Figure 3: SIRS score in pancreatitis subjects**

There was In-hospital mortality rate of 5.2% (n = 3) among 397 cases.[12] Cause of death was found to be Multi-organ Dysfunction Syndrome (MODS). Our study which is comparable to mortality for acute pancreatitis by Singh et al who reported 14 (3.5%) deaths




**Figure 4: Outcome in pancreatitis**

The analysis for prediction of mortality showed higher sensitivity of 100% with BISAP score as compared to 66.7% with PANC3 score. Both scores had equal specificity of 89.09%. Positive and negative predictive values in the study were found to be 25% and 98% respectively for PANC3 score and 33.33% and 100% respectively for BISAP score. Diagnostic accuracy of this study was found to be 84 %, for a PANC3 score of 3 and 97% for BISAP score of 3 or more than 3. We observed that BISAP score was most accurate in predicting mortality (97%). Findings from another study,

doneby Koziel et al. also reported that BISAP was more accurate in predicting mortality when compared to PANC 3 and Ranson's.[13] On the contrary results reported by Park et al[14] and Yadav et al[15] showed better accuracy of the PANC3 score. AUC for BISAP for predicting mortality in AP in our study (0.97) is far better than the study conducted by Park et al[14] (0.86) which might be due to less sample size of our study.

ROC Curve using BISAP and PANC3 score for prediction of



**Figure 5: ROC Curve using BISAP and PANC3 score**

When calculating the likelihood ratios for the BISAP score at a threshold of 3, the positive likelihood ratio was above 5 for both prediction of complications and mortality in pancreatitis subjects, suggesting that a BISAP score of  $\geq 3$  did well in predicting mortality and complications of AP.

Patients were followed for 4 weeks after discharge from the hospital, there was no mortality during this period. 8 patients had AKI at time of discharge. 7 out of them had improvement in renal function tests

## CONCLUSION

BISAP score was found to have more sensitivity, diagnostic accuracy, positive and negative predictive value, and equal specificity in predicting the severity of acute pancreatitis as compared to the PANC3 score. Hence, the BISAP score was found to predict more number of patients with, the likelihood of progressing to severe disease. Larven et al stated in their study that, a prognostic scoring should preferably have high positive predictive values or high negative predictive values to assess the severity of acute pancreatitis. Hence, BISAP is considered a simple and good bedside scoring system in predicting the severity of acute pancreatitis.

The limitations of our study were the small size of study population which limits a more extensive evaluation of the ability of the BISAP and PANC3 scores to predict local complications and mortality. The etiology in this study was found to be different from the worldwide accepted one, hence might not be correct to compare with other studies. Variations in the timing of the presentation of patients to the hospital after the onset of symptoms might have interfered with the assessment of the scoring systems. BISAP score is higher in patients having SIRS, in older patients and in patients with altered mental status. BISAP has the disadvantage that it cannot easily distinguish transient from persistent organ failure.

## REFERENCES

1. Yadav D, Lowenfels AB. The epidemiology of pancreatitis and pancreatic cancer. *Gastroenterology*. 2013 1;144(6):1252-61
2. Mukherjee D, Bhakta S, Lahiry S, Sinha R. Demographic profile of Acute Pancreatitis in Eastern India: A Single Centre Experience. *Asian Journal of Medical Sciences*. 2017;8(6):24-9.
3. Whitcomb DC. Acute pancreatitis. *New England Journal of Medicine*. 2006;354(20):2142-50.
4. Bradley EL 3rd. A clinically based classification system for acute pancreatitis: summary of the International Symposium on Acute Pancreatitis, Atlanta, Ga, September 11–13, 1992. *Arch Surg* 1993;128(5):586–590.
5. Banks PA, Bollen TL, Dervenis C, et al. Classification of acute pancreatitis 2012. Revision of classification and definitions by international consensus. *Gut* 2013;62:102–11.

6. Poppy R, Berger II. Natural II of Acute Paricreautis and the role of/ Infection. *Millers Best Practclingastmenterol* 1999;26(2):286–291.
7. Papachristou GI, Muddana V, Yadav D, O'Connell M, Sanders MK, Slivka A et al. Comparison of BISAP, Ranson's, APACHE-II, and CTSI Scores in Predicting Organ Failure, Complications, and Mortality in Acute Pancreatitis. *American Journal of Gastroenterology*. 2010;105(2):435-41
8. Amodio A, Manfredi R, Katsotourchi AM, et al. Prospective evaluation of subjects with chronic asymptomatic pancreatic hyperenzymemia. *Am J Gastroenterology* 2012;107:1089–95.
9. Lalithkumar J, Chitra T, Kodieswaran N. Comparative study between BISAP and Ranson's score in predicting severity of acute pancreatitis. *Int Arc Intg Med*. 2016;3:23-33.
10. Park JY, Jeon TJ, Ha TH, et al. Bedside index for severity in acute pancreatitis: comparison with other scoring systems in predicting severity and organ failure. *Hepatobiliary Pancreat Dis Int*. 2013;12:645–650
11. A, Rabbani K, Louzi A and Finech B; Interest of SIRS score in acute pancreatitis severity prediction, *International Journal of Surgery Science* 2019; 3(4): 230-231
12. Vikesh K Singh, Bechien U Wu, Thomas L Bollen, Kathryn Repas, Rie Maurer, Koenraad J Morteale, Peter A Banks; Early systemic inflammatory response syndrome is associated with severe acute pancreatitis, *Clin Gastroenterol Hepatol*. 2009 Nov;7(11):1247-51.
13. Koziel D, Gluszek S, Matykiewicz J, Lewitowicz P, Drozdak Z. Comparative analysis of selected scales to assess prognosis in acute pancreatitis. *Canadian journal of gastroenterology & hepatology*. 29(6):299-303.
14. Park JY, Jeon TJ, Ha TH, et al. Bedside index for severity in acute pancreatitis: comparison with other scoring systems in predicting severity and organ failure. *Hepatobiliary Pancreat Dis Int*. 2013;12:645–650.
15. Yadav J, Yadav SK, Kumar S, Baxla RG, Sinha DK, Bodra P, Besra RC, Baski BM, Prakash O, Anand A. Predicting morbidity and mortality in acute pancreatitis in an Indian population: a comparative study of the BISAP score, Ranson's score and CT severity index. *Gastroenterology report*. 2016;4(3):216-20.

**How to cite:** Anubha Garg, Mahima Khatkar, Bharti, Dinesh Kumar Garg, Sumit Chawla. Scoring Systems Using Bedside Index of Severity in Acute Pancreatitis (BISAP) and PANC3 Vs Revised Atlanta Classification in Predicting the Severity of Acute Pancreatitis. *International Medicine*, 2025;11 (1):1-4